arXiv:1211.6978v1 [math.NT] 29 Nov 2012

EXTENDED FERMIONIC p-ADIC q-INTEGRALS ON $\mathbb{Z}_p$ IN CONNECTION WITH APPLICATIONS OF UMBRAL CALCULUS

By

[†]Serkan ARACI, [†]Mehmet ACIKGOZ and [‡]Erdoğan ŞEN

[†]University of Gaziantep, Faculty of Arts and Science, Department of Mathematics, 27310
Gaziantep, Turkey

[‡]Department of Mathematics, Faculty of Science and Letters, Namik Kemal University,
59030 Tekirdağ, Turkey

mtsrkn@hotmail.com; acikgoz@gantep.edu.tr; erdogan.math@gmail.com

Abstract

The purpose of this paper is to derive some applications of umbral calculus by using extended fermionic p-adic q-integral on $\mathbb{Z}_p$. From those applications, we derive some new interesting properties on the new family of Euler numbers and polynomials. That is, a systemic study of the class of Sheffer sequences in connection with generating function of the weighted Euler polynomials are given in the present paper.

2010 Mathematics Subject Classification 11S80, 11B68.

Key Words and Phrases Appell sequence, Sheffer sequence, Euler numbers and polynomials, formal power series, fermionic p-adic q-integral on $\mathbb{Z}_p$.

1. **Preliminaries**

Suppose that p be a fixed odd prime number. Throughout this work we use the following notations, where $\mathbb{Z}_p$ we denote the ring of p-adic rational integers, $\mathbb{Q}$ denotes the field of rational numbers, $\mathbb{Q}_p$ denotes the field of p-adic rational numbers, and $\mathbb{C}_p$ denotes the completion of algebraic closure of $\mathbb{Q}_p$. Let $\mathbb{N}$ be the set of natural numbers and $\mathbb{N}^* = \mathbb{N} \cup \{0\}$. The p-adic absolute value is defined by $|p|_p = p^{-1}$. Also, we assume that $|q-1|_p < 1$ is an indeterminate. Let $UD(\mathbb{Z}_p)$ be the space of uniformly differentiable functions on $\mathbb{Z}_p$. For

$f \in UD(\mathbb{Z}_p)$, the fermionic p-adic q-integral on $\mathbb{Z}_p$ is defined by T. Kim, as follows:

$$I_{-q}(f) = \int_{\mathbb{Z}_p} f(\xi)\, d\mu_{-q}(\xi) = \lim_{n\to\infty} \frac{1}{[p^n]_{-q}} \sum_{\xi=0}^{p^n-1} (-1)^{\xi} f(\xi)\, q^{\xi}. \tag{1.1}$$

where $[x]_q$ is q-analogue of x defined by

$$[x]_q = \frac{q^x - 1}{q-1} \text{ and } [x]_{-q} = \frac{1-(-q)^x}{1+q}.$$

We want to note that $\lim_{q\to 1}[x]_q = x$ (for details, see [1-31]).

By (1.1), we have

$$qI_{-q}(f_1) + I_{-q}(f) = [2]_q f(0) \tag{1.2}$$

where $f_1(\xi) := f(\xi+1)$ (for details, see [3], [8]).

Let us consider Kim's p-adic fermionic q-integral on $\mathbb{Z}_p$ in this form: for $|1-\zeta|_p < 1$

$$I_{-q}^{\zeta}(f) = \int_{\mathbb{Z}_p} \zeta^{\xi} f(\xi)\, d\mu_{-q}(\xi) = \lim_{n\to\infty} \frac{1}{[p^n]_{-q}} \sum_{\xi=0}^{p^n-1} \zeta^{\xi} f(\xi)(-1)^{\xi} q^{\xi}, \tag{1.3}$$

where $I_{-1}^{\zeta}(f)$ are called extended fermionic p-adic q-integral on $\mathbb{Z}_p$.

Let us now consider $f_1(\xi) := f(\xi+1)$, then we develop as follows:

$$\begin{aligned} -q\zeta I_{-q}^{\zeta}(f_1) &= \lim_{n\to\infty} \frac{1}{[p^n]_{-q}} \sum_{\xi=0}^{p^n-1} \zeta^{\xi+1}(-1)^{\xi+1} f(\xi+1)\, q^{\xi+1} \\ &= I_{-q}^{\zeta}(f) + \frac{[2]_q}{2} \lim_{n\to\infty} \left(-f(0) - \xi^{p^n} f(p^n)\, q^{p^n}\right) \\ &= I_{-q}^{\zeta}(f) - [2]_q f(0). \end{aligned}$$

Therefore, we have the following lemma.

Lemma 1. *For $f \in UD(\mathbb{Z}_p)$, we get*

$$q\zeta I_{-q}^{\zeta}(f_1) + I_{-q}^{\zeta}(f) = [2]_q f(0).$$

Taking $f(\xi) = e^{t(x+\xi)} \in UD(\mathbb{Z}_p)$ in Lemma 1, then we introduce the following expression:

$$\int_{\mathbb{Z}_p} \zeta^{\xi} e^{t(x+\xi)} d\mu_{-q}(\xi) = \frac{[2]_q}{q\zeta e^t + 1} e^{tx} = \sum_{n=0}^{\infty} E_{n,\zeta}^{q}(x) \frac{t^n}{n!}, \tag{1.4}$$

where we call $E_{n,\zeta}^{q}(x)$ as weighted q-Euler polynomials. In the special case, $x = 0$, $E_{n,\zeta}^{q}(0) := E_{n,\zeta}^{q}$ are called weighted q-Euler numbers and the relation between weighted q-Euler numbers and weighted q-Euler polynomials are given by

$$E_{n,\zeta}^{q}(x) = \sum_{l=0}^{n} \binom{n}{l} x^l E_{n-l,\zeta}^{q} = \left(x + E_{\zeta}^{q}\right)^n, \tag{1.5}$$

with the usual of replacing $\left(E_{\zeta}^{q}\right)^{n}$ by $E_{n,\zeta}^{q}$ is used. By (1.4), we note that

$$E_{n,\zeta}^{q}=\int_{\mathbb{Z}_p}\zeta^{\xi}\xi^{n}d\mu_{-q}\left(\xi\right)\ \ and\ E_{n,\zeta}^{q}\left(x\right)=\int_{\mathbb{Z}_p}\zeta^{\xi}\left(x+\xi\right)^{n}d\mu_{-q}\left(\xi\right). \tag{1.6}$$

By (1.4), we have

$$E_{n,\zeta}^{q}\left(x\right)=\left[2\right]_{q}\sum_{m=0}^{\infty}\left(-1\right)^{m}q^{m}\zeta^{m}\left(m+x\right)^{n},\ for\ n\in\mathbb{N}^{*}. \tag{1.7}$$

From this, we can define weighted q-Zeta function as follows:

$$\lambda\left(s,x:q:\zeta\right)=\left[2\right]_{q}\sum_{m=0}^{\infty}\frac{\left(-1\right)^{m}q^{m}\zeta^{m}}{\left(m+x\right)^{s}} \tag{1.8}$$

By (1.7) and (1.8), we derive the following equation (1.9):

$$\lambda\left(-n,x:q:\zeta\right)=E_{n,\zeta}^{q}\left(x\right),\ for\ any\ n\in\mathbb{N}^{*}. \tag{1.9}$$

When we set as $q=\zeta=1$ in (1.9) which reduces to

$$\zeta_{E}\left(-n,x\right)=E_{n}\left(x\right)$$

which is well known in [11].

By (1.3) and (1.4), we compute

$$\begin{aligned}\int_{\mathbb{Z}_p}\zeta^{\xi}\left(x+\xi\right)^{n}d\mu_{-q}\left(\xi\right) &= \lim_{m\rightarrow\infty}\frac{1}{\left[dp^{m}\right]_{-q}}\sum_{\xi=0}^{dp^{m}-1}\left(-1\right)^{\xi}\zeta^{\xi}\left(x+\xi\right)^{n}q^{\xi}\\ &= \frac{d^{n}}{\left[d\right]_{-q}}\sum_{j=0}^{d-1}\left(-1\right)^{j}\zeta^{j}q^{j}\left(\lim_{m\rightarrow\infty}\frac{1}{\left[p^{m}\right]_{\left(-q\right)^{d}}}\sum_{\xi=0}^{p^{m}-1}\left(-1\right)^{\xi}\left(\zeta^{d}\right)^{\xi}\left(q^{d}\right)^{\xi}\left(\frac{x+j}{d}+\xi\right)^{n}\right)\\ &= \frac{d^{n}}{\left[d\right]_{-q}}\sum_{j=0}^{d-1}\left(-1\right)^{j}\zeta^{j}q^{j}\int_{\mathbb{Z}_p}\zeta^{d\xi}\left(\frac{x+j}{d}+\xi\right)^{n}d\mu_{-q^{d}}\left(\xi\right),\end{aligned}$$

where d is an odd natural number. So from the above

$$\int_{\mathbb{Z}_p}\zeta^{\xi}\left(x+\xi\right)^{n}d\mu_{-q}\left(\xi\right)=\frac{d^{n}}{\left[d\right]_{-q}}\sum_{j=0}^{d-1}\left(-1\right)^{j}\zeta^{j}q^{j}\int_{\mathbb{Z}_p}\zeta^{d\xi}\left(\frac{x+j}{d}+\xi\right)^{n}d\mu_{-q^{d}}\left(\xi\right). \tag{1.10}$$

By (1.6) and (1.10), we get

$$E_{n,\zeta}^{q}\left(dx\right)=\frac{d^{n}}{\left[d\right]_{-q}}\sum_{j=0}^{d-1}\left(-1\right)^{j}\zeta^{j}q^{j}E_{n,\zeta^{d}}^{q^{d}}\left(x+\frac{j}{d}\right), \tag{1.11}$$

which plays an important role for studying regarding Measure theory on p-adic analysis.

Let we use the following notations, where $\mathbb{C}$ denotes the set of complex numbers, $\mathcal{F}$ denotes the set of all formal power series in the variable t over $\mathbb{C}$ with

$$\mathcal{F}=\left\{f\left(t\right)=\sum_{k=0}^{\infty}a_{k}\frac{t^{k}}{k!}\mid a_{k}\in\mathbb{C}\right\},$$

$\mathcal{P}=\mathbb{C}[x]$ and $\mathcal{P}^*$ denotes the vector space of all linear functional on $\mathcal{P}$, $\langle L \mid p(x)\rangle$ denotes the action of the linear functional L on the polynomial $p(x)$, and it is well-known that the vector space operation on $\mathcal{P}^*$ is defined by

$$\begin{aligned} \langle L+M \mid p(x)\rangle &= \langle L \mid p(x)\rangle + \langle M \mid p(x)\rangle, \\ \langle cL \mid p(x)\rangle &= c\langle L \mid p(x)\rangle, \end{aligned}$$

where c is any constant in $\mathbb{C}$ (for details, see [12], [19], [13], [14], [31]).

The formal power series are known by

$$f(t)=\sum_{k=0}^{\infty} a_k \frac{t^k}{k!} \in \mathcal{F}$$

which describes a linear functional on $\mathcal{P}$ as $\langle f(t) \mid x^n\rangle = a_n$ for all $n \geq 0$ (for details, see [12], [19], [13], [14], [31]). In addition to

$$\left\langle t^k \mid x^n \right\rangle = n!\delta_{n,k}, \tag{1.12}$$

where $\delta_{n,k}$ is the Kronecker delta. If we take as

$$f_L(t)=\sum_{k=0}^{\infty}\left\langle L \mid x^k\right\rangle \frac{t^k}{k!},$$

then we obtain

$$\langle f_L(t) \mid x^n\rangle = \langle L \mid x^n\rangle$$

and so as linear functionals $L=f_L(t)$ (see [12], [19], [13], [14], [31]). Additionally, the map $L \rightarrow f_L(t)$ is a vector space isomorphism from $\mathcal{P}^*$ onto $\mathcal{F}$. Henceforth, $\mathcal{F}$ will denote both the algebra of the formal power series in t and the vector space of all linear functionals on $\mathcal{P}$, and so an element $f(t)$ of $\mathcal{F}$ will be thought of as both a formal power series and a linear functional. $\mathcal{F}$ will be called as umbral algebra (see [12], [19], [13], [14], [31]).

It is well-known that $\langle e^{yt} \mid x^n\rangle = y^n$. Then, leads to the following

$$\left\langle e^{yt} \mid p(x)\right\rangle = p(y)$$

(see [12], [19], [13], [14], [15], [31]). We want to note that for all $f(t)$ in $\mathcal{F}$

$$f(t)=\sum_{k=0}^{\infty}\left\langle f(t) \mid x^k\right\rangle \frac{t^k}{k!} \tag{1.13}$$

and for all polynomial $p(x)$,

$$p(x)=\sum_{k=0}^{\infty}\left\langle t^k \mid p(x)\right\rangle \frac{x^k}{k!}, \tag{1.14}$$

(for details, see [12], [19], [13], [14], [31]). The order $o(f(t))$ of the power series $f(t)\neq 0$ is the smallest integer k for which a_k does not vanish. It is considered $o(f(t))=\infty$ if $f(t)=0$. We see that $o(f(t)g(t)) = o(f(t)) + o(g(t))$ and $o(f(t)+g(t)) \geq \min\{o(f(t)), o(g(t))\}$. The series $f(t)$ has a multiplicative inverse, denoted by $f(t)^{-1}$ or $\frac{1}{f(t)}$, if and only if $o(f(t))=0$. Such series is called an invertible series. A series $f(t)$ for which $o(f(t))=1$ is called a delta series (see [12], [19], [13], [14], [15], [31]). For $f(t), g(t) \in \mathcal{F}$, we have $\langle f(t)g(t) \mid p(x)\rangle = \langle f(t) \mid g(t)p(x)\rangle$.

A delta series $f(t)$ has a compositional inverse $\overline{f}(t)$ such that $f\left(\overline{f}(t)\right)=\overline{f}(f(t))=t$. For $f(t), g(t)\in\mathcal{F}$, we have $\langle f(t)g(t)\mid p(x)\rangle=\langle f(t)\mid g(t)p(x)\rangle$. By (1.13), we have

$$p^{(k)}(x)=\frac{d^kp(x)}{dx^k}=\sum_{l=k}^{\infty}\frac{\left\langle t^l\mid p(x)\right\rangle}{l!}l(l-1)\cdots(l-k+1)x^{l-k}. \tag{1.15}$$

Thus, notice that

$$p^{(k)}(0)=\left\langle t^k\mid p(x)\right\rangle=\left\langle 1\mid p^{(k)}(x)\right\rangle. \tag{1.16}$$

By (1.15), we have

$$t^kp(x)=p^{(k)}(x)=\frac{d^kp(x)}{dx^k}. \tag{1.17}$$

So from the above

$$e^{yt}p(x)=p(x+y). \tag{1.18}$$

Let $S_n(x)$ be a polynomial with $\deg S_n(x)=n$. Let $f(t)$ be a delta series and let $g(t)$ be an invertible series. Then there exists a unique sequence $S_n(x)$ of polynomials such that $\left\langle g(t)f(t)^k\mid S_n(x)\right\rangle=n!\delta_{n,k}$ for all $n,k\geq 0$. The sequence $S_n(x)$ is called the Sheffer sequence for $(g(t), f(t))$ or that $S_n(t)$ is Sheffer for $(g(t), f(t))$.

The Sheffer sequence for $(1, f(t))$ is called the associated sequence for $f(t)$ or $S_n(x)$ is associated to $f(t)$. The sheffer sequence for $(g(t), t)$ is called the Appell sequence for $g(t)$ or $S_n(x)$ is Appell for $g(t)$.

Let $p(x)\in\mathcal{P}$. Then we have

$$\begin{aligned}\langle f(t)\mid xp(x)\rangle &= \langle\partial_t f(t)\mid p(x)\rangle=\langle f'(t)\mid p(x)\rangle, \\ \langle e^y+1\mid p(x)\rangle &= p(y)+p(0),\ \text{(see [31])}.\end{aligned} \tag{1.19}$$

Let $S_n(x)$ be sheffer for $(g(t), f(t))$. Then

$$\begin{aligned}h(t) &= \sum_{k=0}^{\infty}\frac{\langle h(t)\mid S_k(x)\rangle}{k!}g(t)f(t)^k,\ h(t)\in\mathcal{F} \\ p(x) &= \sum_{k=0}^{\infty}\frac{\left\langle g(t)f(t)^k\mid p(x)\right\rangle}{k!}S_k(x),\ p(x)\in\mathcal{P}, \\ \frac{1}{g\left(\overline{f}(t)\right)}e^{y\overline{f}(t)} &= \sum_{k=0}^{\infty}S_k(y)\frac{t^k}{k!},\ \text{for all } y\in\mathbb{C}, \\ f(t)S_n(x) &= nS_{n-1}(x).\end{aligned} \tag{1.20}$$

Also, it is well known in [31] that

$$\langle f_1(t)f_2(t)\cdots f_m(t)\mid x^n\rangle=\sum\binom{n}{i_1,\cdots,i_m}\left\langle f_1(t)\mid x^{i_1}\right\rangle\cdots\left\langle f_m(t)\mid x^{i_m}\right\rangle \tag{1.21}$$

where $f_1(t), f_2(t),\cdots, f_m(t)\in\mathcal{F}$ and the sum is over all nonnegative integers $i_1,\cdots,i_m$ such that $i_1+\cdots+i_m=n$ (see [31]).

In [19] and [20], Dere and Simsek have studied applications of umbral algebra to special functions. They gave some new interesting links for further works of many mathematicians in Analytic number theory and in modern classical umbral calculus. Kim *et al.* have given some properties of umbral calculus for Frobenius-Euler polynomials [12], Euler polynomials [13] and other special functions [14]. Also, they investigated some new applications of umbral calculus associated with p-adic invariants integral on $\mathbb{Z}_p$ in [1].

By the same motivation, we also give some applications of umbral calculus by using extended fermionic p-adic integral on $\mathbb{Z}_p$. From those applications, we derive some interesting equalities on weighted Euler numbers, weighted Euler polynomials and weighted Euler polynomials of order k.

2. On the extended fermionic p-adic q-integral on $\mathbb{Z}_p$ in connection with applications of umbral calculus

Suppose that $S_n(x)$ is an Appell sequence for $g(t)$. Then, by (1.20), we have

$$\frac{1}{g(t)}x^n = S_n(x) \text{ if and only if } x^n = g(t)\,S_n(x)\,, \ (n \geq 0)\,. \tag{2.1}$$

Let us contemplate as follows:

$$g_q(t \mid \zeta) = \frac{q\zeta e^t + 1}{[2]_q} \in \mathcal{F}.$$

Therefore, we easily notice that $g(t)$ is an invertible series. By (2.1), we have

$$\sum_{n=0}^{\infty} E_{n,\zeta}^{q}(x)\frac{t^n}{n!} = \frac{1}{g_q(t \mid \zeta)}e^{xt}. \tag{2.2}$$

By (2.2), we have

$$\frac{1}{g_q(t \mid \zeta)}x^n = E_{n,\zeta}^{q}(x)\,. \tag{2.3}$$

Also, by (1.20), we have

$$tE_{n,\zeta}^{q}(x) = \left(E_{n,\zeta}^{q}(x)\right)^{\prime} = nE_{n-1,\zeta}^{q}(x)\,, \tag{2.4}$$

By (2.3) and (2.4), we have the following proposition.

Proposition 1. *For $n \geq 0$, $E_{n,\zeta}^{q}(x)$ is an Appell sequence for $g_q(t \mid \zeta) = \frac{\zeta q e^t + 1}{[2]_q}$.*

By (1.6), we derive that

$$\begin{aligned}\sum_{n=1}^{\infty} E_{n,\zeta}^{q}(x)\frac{t^n}{n!} &= \frac{xg_q(t \mid \zeta)\,e^{xt} - \acute{g}_q(t \mid \zeta)\,e^{xt}}{g(t)^2} \\ &= \sum_{n=0}^{\infty}\left(x\frac{1}{g_q(t \mid \zeta)}x^n - \frac{\acute{g}_q(t \mid \zeta)}{g_q(t \mid \zeta)}\frac{1}{g_q(t \mid \zeta)}x^n\right)\frac{t^n}{n!}\end{aligned} \tag{2.5}$$

Because of (2.3) and (2.5), we discover the following:

$$E_{n+1,\zeta}^{q}(x) = xE_{n,\zeta}^{q}(x) - \frac{\acute{g}_q(t \mid \zeta)}{g_q(t \mid \zeta)}E_{n,\zeta}^{q}(x)\,.$$

Therefore, we get the following theorem.

Theorem 1. *Let* $g_q(t \mid \zeta) = \frac{\zeta q e^t + 1}{2} \in \mathcal{F}$. *Then we have for* $n \geq 0$:

$$E^q_{n+1,\zeta}(x) = \left(x - \frac{g\acute{}_q(t \mid \zeta)}{g_q(t \mid \zeta)}\right) E^q_{n,\zeta}(x). \tag{2.6}$$

Moreover,

$$\lambda(-n-1, x : q : \zeta) = \left(x - \frac{g\acute{}_q(t \mid \zeta)}{g_q(t \mid \zeta)}\right) \lambda(-n, x : q : \zeta).$$

where $g\acute{}_q(t \mid \zeta) = \frac{dg_q(t|\zeta)}{dt}$.

From (1.6), it is easy to show that

$$\sum_{n=0}^{\infty} \left(\zeta q E^q_{n,\zeta}(x+1) + E^q_{n,\zeta}(x)\right) \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left([2]_q x^n\right) \frac{t^n}{n!}.$$

By comparing the coefficients in the both sides of $\frac{t^n}{n!}$ on the above, we develop the following:

$$\zeta q E^q_{n,\zeta}(x+1) + E^q_{n,\zeta}(x) = [2]_q x^n. \tag{2.7}$$

From Theorem 1, we get the following equation (2.8):

$$g_q(t \mid \zeta) E^q_{n+1,\zeta}(x) = g_q(t \mid \zeta) x E^q_{n,\zeta}(x) - g\acute{}_q(t \mid \zeta) E^q_{n,\zeta}(x). \tag{2.8}$$

So from above

$$\left(\zeta q e^t + 1\right) E^q_{n+1,\zeta}(x) = \left(\zeta q e^t + 1\right) x E^q_{n,\zeta}(x) - \zeta q e^t E^q_{n,\zeta}(x).$$

Thus, we can write the following equation:

$$\zeta q E^q_{n+1,\zeta}(x+1) + E^q_{n+1,\zeta}(x) = \zeta q (x+1) E^q_{n,\zeta}(x+1) + x E^q_{n,\zeta}(x) - \zeta q E^q_{n,\zeta}(x+1). \tag{2.9}$$

From (2.7) (2.8) and (2.9), we can state following theorem.

Theorem 2. *For* $n \geq 0$, *then we have*

$$\zeta q E^q_{n,\zeta}(x+1) + E^q_{n,\zeta}(x) = [2]_q x^n. \tag{2.10}$$

Remark 1. *Assume that* $S_n(x)$ *is Sheffer sequence for* $(g(t), f(t))$. *Then Sheffer identity is introduced by*

$$S_n(x+y) = \sum_{k=0}^{n} \binom{n}{k} P_k(y) S_{n-k}(x) = \sum_{k=0}^{n} \binom{n}{k} P_k(x) S_{n-k}(y), \tag{2.11}$$

where $P_k(y) = S_k(y) g(t)$ *is associated to* $f(t)$ *(for details, see* [13], [19], [20], [31]*).*

On account of (1.4) and (2.11), then we have

$$\begin{aligned} E^q_{n,\zeta}(x+y) &= \sum_{k=0}^{n} \binom{n}{k} P_k(y) S_{n-k}(x) \\ &= \sum_{k=0}^{n} \binom{n}{k} E^q_{n-k,\zeta}(y) x^k. \end{aligned}$$

So we have

$$E_{n,\zeta}^{q}\left(x+y\right)=\sum_{k=0}^{n}\binom{n}{k}E_{n-k,\zeta}^{q}\left(y\right)x^{k}.$$

By (1.4), we easily see for $\alpha\left(\neq 0\right)\in\mathbb{C}$:

$$E_{n,\zeta}^{q}\left(\alpha x\right)=\frac{g_{q}\left(t\mid\zeta\right)}{g_{q}\left(\frac{t}{\alpha}\mid\zeta\right)}E_{n,\zeta}^{q}\left(x\right). \tag{2.12}$$

From (1.11) and (2.12), we readily derive for $d\equiv 1\left(\operatorname{mod}2\right)$:

$$\frac{g_{q}\left(t\mid\zeta\right)}{g_{q}\left(\frac{t}{d}\mid\zeta\right)}E_{n,\zeta}^{q}\left(x\right)=\frac{d^{n}}{\left[d\right]_{-q}}\sum_{j=0}^{d-1}\left(-1\right)^{j}\zeta^{j}q^{j}E_{n,\zeta^{d}}^{q^{d}}\left(x+\frac{j}{d}\right).$$

Let us consider the linear functional $f\left(t\right)$ that satisfies:

$$\left\langle f\left(t\right)\mid p\left(x\right)\right\rangle=\int_{\mathbb{Z}_{p}}\zeta^{\xi}p\left(\xi\right)d\mu_{-q}\left(\xi\right), \tag{2.13}$$

for all polynomials $p\left(x\right)$. From (2.13), we readily see that

$$f\left(t\right)=\sum_{n=0}^{\infty}\frac{\left\langle f\left(t\right)\mid x^{n}\right\rangle}{n!}t^{n}=\sum_{n=1}^{\infty}\left(\int_{\mathbb{Z}_{p}}\zeta^{\xi}\xi^{n}d\mu_{-q}\left(\xi\right)\right)\frac{t^{n}}{n!}=\int_{\mathbb{Z}_{p}}\zeta^{\xi}e^{\xi t}d\mu_{-q}\left(\xi\right). \tag{2.14}$$

Thus, we have

$$f\left(t\right)=\int_{\mathbb{Z}_{p}}\zeta^{\xi}e^{\xi t}d\mu_{-q}\left(\xi\right)=\frac{\left[2\right]_{q}}{\zeta qe^{t}+1}. \tag{2.15}$$

Therefore, by (2.13) and (2.15), we arrive at the following theorem.

Theorem 3. *For $n\geq 0$, then we have*

$$\left\langle f\left(t\right)\mid p\left(x\right)\right\rangle=\int_{\mathbb{Z}_{p}}\zeta^{\xi}p\left(\xi\right)d\mu_{-q}\left(\xi\right). \tag{2.16}$$

Also,

$$\left\langle\frac{\left[2\right]_{q}}{\zeta qe^{t}+1}\mid p\left(x\right)\right\rangle=\int_{\mathbb{Z}_{p}}\zeta^{\xi}p\left(\xi\right)d\mu_{-q}\left(\xi\right). \tag{2.17}$$

Obviously that

$$E_{n,\zeta}^{q}=\left\langle\int_{\mathbb{Z}_{p}}\zeta^{\xi}e^{\xi t}d\mu_{-q}\left(\xi\right)\mid x^{n}\right\rangle. \tag{2.18}$$

From (1.6) and (2.18), we see that

$$\sum_{n=0}^{\infty}\left(\int_{\mathbb{Z}_{p}}\zeta^{\xi}\left(x+\xi\right)^{n}d\mu_{-q}\left(\xi\right)\right)\frac{t^{n}}{n!}=\int_{\mathbb{Z}_{p}}\zeta^{\xi}e^{\left(x+\xi\right)t}d\mu_{-q}\left(\xi\right)=\sum_{n=0}^{\infty}\left(\int_{\mathbb{Z}_{p}}\zeta^{\xi}e^{\xi t}d\mu_{-q}\left(\xi\right)x^{n}\right)\frac{t^{n}}{n!}. \tag{2.19}$$

By (1.6) and (2.20), we see that for $n\in\mathbb{N}^{*}$:

$$E_{n,\zeta}^{q}\left(x\right)=\int_{\mathbb{Z}_{p}}\zeta^{\xi}\left(x+\xi\right)^{n}d\mu_{-q}\left(\xi\right)=\int_{\mathbb{Z}_{p}}\zeta^{\xi}e^{\xi t}d\mu_{-q}\left(\xi\right)x^{n}. \tag{2.20}$$

Consequently, we obtain the following theorem.

Theorem 4. *For* $p(x) \in \mathcal{P}$, *then we have*

$$\begin{aligned} \int_{\mathbb{Z}_p} \zeta^{\xi} p(x+\xi)\, d\mu_{-q}(\xi) &= \int_{\mathbb{Z}_p} \zeta^{\xi} e^{\xi t} d\mu_{-q}(\xi)\, p(x) \\ &= \frac{[2]_q}{\zeta q e^t + 1} p(x). \end{aligned} \tag{2.21}$$

That is:

$$E_{n,\zeta}^{q}(x) = \int_{\mathbb{Z}_p} \zeta^{\xi} e^{\xi t} d\mu_{-q}(\xi)\, x^n = \frac{[2]_q}{\zeta q e^t + 1} x^n. \tag{2.22}$$

For $|1-\zeta|_p < 1$, we introduce weighted q-Euler polynomials of order k as follows:

$$\begin{aligned} \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{\text{k-times}} \zeta^{\xi_1+\cdots+\xi_k} e^{(\xi_1+\cdots+\xi_k+x)t} d\mu_{-q}(\xi_1)\cdots d\mu_{-q}(\xi_k) &= \left(\frac{[2]_q}{q\zeta e^t+1}\right)^k e^{xt} \\ &= \sum_{n=0}^{\infty} E_{n,\zeta}^{(k)}(x \mid q) \frac{t^n}{n!}. \end{aligned} \tag{2.23}$$

where, for $x=0$, $E_{n,\zeta}^{(k)}(0 \mid q) := E_{n,\zeta}^{(k)}(q)$ are called weighted q-Euler numbers of order k.

By (2.23), we have

$$\begin{aligned} &\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{\text{k-times}} \zeta^{\xi_1+\cdots+\xi_k} (\xi_1+\cdots+\xi_k+x)^n d\mu_{-q}(\xi_1)\cdots d\mu_{-q}(\xi_k) \\ = &\sum_{i_1+\cdots+i_k=n} \binom{n}{i_1,\cdots,i_m} \int_{\mathbb{Z}_p} \zeta^{\xi_1}\xi_1^{i_1} d\mu_{-q}(\xi_1)\cdots \int_{\mathbb{Z}_p} \zeta^{\xi_k}\xi_k^{i_k} d\mu_{-q}(\xi_k) \\ = &\sum_{i_1+\cdots+i_k=n} \binom{n}{i_1,\cdots,i_m} E_{i_1,\zeta}^{q}\cdots E_{i_k,\zeta}^{q} = E_{n,\zeta}^{(k)}(x \mid q). \end{aligned} \tag{2.24}$$

Thanks to (2.23) and (2.24), we have

$$E_{n,\zeta}^{(k)}(x \mid q) = \sum_{l=0}^{n} \binom{n}{l} x^l E_{n-l,\zeta}^{(k)}(q). \tag{2.25}$$

From (2.24) and (2.25), we notice that $E_{n,\zeta}^{(k)}(x \mid q)$ is a monic polynomial of degree n with coefficients in $\mathbb{Q}$. For $k \in \mathbb{N}$, let us assume that

$$\begin{aligned} g_q^{(k)}(t \mid \zeta) &= \left(\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{\text{k-times}} \zeta^{\xi_1+\cdots+\xi_k} e^{(\xi_1+\cdots+\xi_k)t} d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) \right)^{-1} \\ &= \left(\frac{\zeta q e^t + 1}{[2]_q} \right)^k . \end{aligned} \tag{2.26}$$

From (2.26), we note that $g_q^{(k)}(t \mid \zeta)$ is an invertible series. On account of (2.23) and (2.26), we readily derive that

$$\begin{aligned} \frac{1}{g_q^{(k)}(t \mid \zeta)} e^{xt} &= \underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{\text{k-times}} \zeta^{\xi_1+\cdots+\xi_k} e^{(\xi_1+\cdots+\xi_k+x)t} d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) \\ &= \sum_{n=0}^{\infty} E_{n,\zeta}^{(k)}(x \mid q) \frac{t^n}{n!}. \end{aligned} \tag{2.27}$$

Also, we note that

$$tE_{n,\zeta}^{(k)}(x \mid q) = nE_{n-1,\zeta}^{(k)}(x \mid q). \tag{2.28}$$

By (2.27) and (2.28), we easily see that $E_{n,\zeta}^{(k)}(x \mid q)$ is an Appell sequence for $g_q^{(k)}(t \mid \zeta)$. Then, by (2.27) and (2.28), we get the following theorem.

Theorem 5. *For $p(x) \in \mathcal{P}$ and $k \in \mathbb{N}$, we have*

$$\underbrace{\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p}}_{\textit{k-times}} \zeta^{\xi_1+\cdots+\xi_k} p(\xi_1+\cdots+\xi_k+x) d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) = \left(\frac{[2]_q}{\zeta q e^t + 1} \right)^k p(x). \tag{2.29}$$

In the special case, the weighted q-Euler polynomials of degree k are derived by

$$E_{n,\zeta}^{(k)}(x \mid q) = \left(\frac{2}{\zeta q e^t + 1} \right)^k x^n = \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \zeta^{\xi_1+\cdots+\xi_k} e^{(\xi_1+\cdots+\xi_k)t} d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) x^n$$

Thus, we get

$$E_{n,\zeta}^{(k)}(x \mid q) \sim \left(\left(\frac{\zeta q e^t + 1}{[2]_q} \right)^k, t \right).$$

Let us take the linear functional $f^{(k)}(t)$ that satisfies

$$\left\langle f^{(k)}(t) \mid p(x) \right\rangle = \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \zeta^{\xi_1+\cdots+\xi_k} p(\xi_1+\cdots+\xi_k) d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k), \tag{2.30}$$

for all polynomials $p(x)$. Therefore, we compute as follows:

$$\begin{aligned}
f^{(k)}(t) &= \sum_{n=0}^{\infty} \frac{\left\langle f^{(k)}(t) \mid x^n \right\rangle}{n!} t^n \\
&= \sum_{n=0}^{\infty} \left(\int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \zeta^{\xi_1+\cdots+\xi_k} \left(\xi_1+\cdots+\xi_k\right)^n d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) \right) \frac{t^n}{n!} \\
&= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \zeta^{\xi_1+\cdots+\xi_k} e^{(\xi_1+\cdots+\xi_k)t} d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) \\
&= \left(\frac{[2]_q}{\zeta q e^t + 1} \right)^k .
\end{aligned}$$

Therefore, the following theorem can be expressed.

Theorem 6. *For $p(x) \in \mathcal{P}$, we have*

$$\begin{aligned}
&\left\langle \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \zeta^{\xi_1+\cdots+\xi_k} e^{(\xi_1+\cdots+\xi_k)t} d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) \mid p(x) \right\rangle \\
&= \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \zeta^{\xi_1+\cdots+\xi_k} p(\xi_1+\cdots+\xi_k) d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k)
\end{aligned}$$

Furthermore,

$$\left\langle \left(\frac{[2]_q}{\zeta q e^t + 1} \right)^k \mid p(x) \right\rangle = \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \zeta^{\xi_1+\cdots+\xi_k} p(\xi_1+\cdots+\xi_k) d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) .$$

That is:

$$E_{n,\zeta}^{(k)}(q) = \left\langle \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \zeta^{\xi_1+\cdots+\xi_k} e^{(\xi_1+\cdots+\xi_k)t} d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) \mid x^n \right\rangle .$$

From (1.21), we notice that

$$\begin{aligned}
&\left\langle \int_{\mathbb{Z}_p} \cdots \int_{\mathbb{Z}_p} \zeta^{\xi_1+\cdots+\xi_k} e^{(\xi_1+\cdots+\xi_k)t} d\mu_{-q}(\xi_1) \cdots d\mu_{-q}(\xi_k) \mid x^n \right\rangle \\
&= \sum_{i_1+\cdots+i_k=n} \binom{n}{i_1, \cdots, i_m} \left\langle \int_{\mathbb{Z}_p} \zeta^{\xi_1} e^{\xi_1 t} d\mu_{-q}(\xi_1) \mid x^{i_1} \right\rangle \cdots \left\langle \int_{\mathbb{Z}_p} \zeta^{\xi_k} e^{\xi_k t} d\mu_{-q}(\xi_k) \mid x^{i_k} \right\rangle
\end{aligned}$$

Therefore, we have

$$E_{n,\zeta}^{(k)}(q) = \sum_{i_1+\cdots+i_k=n} \binom{n}{i_1, \cdots, i_m} E_{i_1,\zeta}^{q} \cdots E_{i_k,\zeta}^{q}.$$

Remark 2. *Our applications for weighted Euler polynomials, weighted q-Euler numbers and weighted q-Euler polynomials of order k seem to be interesting for evaluating at $w = \zeta = 1$*

which lead to Euler polynomials and Euler polynomials of order k, are defined respectively by

$$\begin{aligned}\sum_{n=0}^{\infty} E_n(x)\frac{t^n}{n!} &= \frac{2}{e^t+1}e^{xt},\\ \sum_{n=0}^{\infty} E_n^{(k)}(x)\frac{t^n}{n!} &= \left(\frac{2}{e^t+1}\right)^k e^{xt}.\end{aligned}$$

Also, it is well known that they have representations in terms of fermionic p-adic integral on $\mathbb{Z}_p$ as follows:

$$\begin{aligned}E_n(x) &= \int_{\mathbb{Z}_p}(x+\xi)^n\,d\mu_{-1}(\xi),\\ E_n^{(k)}(x) &= \int_{\mathbb{Z}_p}\cdots\int_{\mathbb{Z}_p}(\xi_1+\cdots+\xi_k+x)^n\,d\mu_{-1}(\xi_1)\cdots d\mu_{-1}(\xi_k).\end{aligned}$$

References

[1] T. Kim and D. S. Kim, Applications of umbral calculus associated with p-adic invariants integral on $\mathbb{Z}_p$, *Abstract and Applied Analysis*, Vol. 2012 (**2012**), Article ID 865721, pages 12 (In press).

[2] T. Kim, D. S. Kim, S. Jeong and S. H. Rim, A note on Fermionic p-adic integrals on $\mathbb{Z}_p$ and umbral calculus, arXiv:1211.6638 [math.NT].

[3] T. Kim, Symmetry p-adic invariant integral on $\mathbb{Z}_p$ for Bernoulli and Euler polynomials, *J. Difference Equ. Appl.* 14 (**2008**), no. 12, 1267-1277.

[4] T. Kim, The modified q-Euler numbers and polynomials, *Adv. Stud. Contemp. Math.* 16 (**2008**), 161–170.

[5] T. Kim, p-adic q-integrals associated with the Changhee–Barnes' q-Bernoulli polynomials, *Integral Transforms Spec. Funct.* 15 (**2004**) 415–420

[6] T. Kim, On p-adic interpolating function for q-Euler numbers and its derivatives, *J. Math. Anal. Appl.* 339 (**2008**), 598–608.

[7] T. Kim, On a q-analogue of the p-adic log gamma functions and related integrals, *J. Number Theory* 76 (**1999**), 320-329.

[8] T. Kim, Symmetry of power sum polynomials and multivariate fermionic p-adic invariant integral on $\mathbb{Z}_p$, *Russ. J. Math. Phys.* 16, no. 1, 93-96 (**2009**).

[9] T. Kim, On the q-extension of Euler and Genocchi numbers, J. Math. Anal. Appl. 326 (**2007**) 1458–1465.

[10] T. Kim, Non-Archimedean q-integrals associated with multiple Changhee q-Bernoulli polynomials, *Russ. J. Math. Phys.* 10 (**2003**) 91–98.

[11] T. Kim, Euler numbers and polynomials associated with zeta functions, *Abstract and Applied Analysis*, Volume **2008**, Article ID 581582, 11 pages.

[12] D. S. Kim and T. Kim, Some identities of Frobenius-Euler polynomials arising from umbral calculus, *Advances in Difference Equations* (In press).

[13] D. S. Kim, T. Kim and S. H. Rim, Umbral calculus and Euler polynomials, arXiv:1211.6639 [math.NT].

[14] D. S. Kim, T. Kim and S. H. Rim, Some identities of polynomials arising from umbral calculus, arXiv:1211.3738 [math.NT].

[15] M. Maldonado, J. Prada and M. J. Senosiain, Appell bases on sequence spaces, *Journal of Nonlinear Mathematical Physics,* Vol. 18, Suppl. 1 (**2011**) 189–194.

[16] M. Acikgoz and Y. Simsek, On multiple interpolation function of the Nörlund-type q-Euler polynomials, *Abst. Appl. Anal.* 2009 (**2009**), Article ID 382574, 14 pages.

[17] M. Acikgoz, S. Araci and I. N. Cangul, A note on the modified q-Bernstein polynomials for functions of several variables and their reflections on q-Volkenborn integration, *Applied Mathematics and Computation*, vol. 218, no. 3, pp. 707–712, **2011**.

[18] I. N. Cangul, V. Kurt, H. Ozden and Y. Simsek, On the higher-order w-q-Genocchi numbers, *Adv. Stud. Contemp. Math.* 19 (**2009**), no. 1, 39–57.

[19] R. Dere and Y. Simsek, Applications of umbral algebra to some special polynomials, *Adv. Stud. Contemp. Math.* 22 (**2012**), no. 3, 433-438.

[20] R. Dere and Y. Simsek, Genocchi polynomials associated with the Umbral algebra, *Applied Mathematics and Computation,* vol. 218, no. 3, pp. 756-761, **2011**

[21] S. Araci, D. Erdal and J. J. Seo, A study on the fermionic p-adic q-integral representation on $\mathbb{Z}_p$ associated with weighted q-Bernstein and q-Genocchi polynomials, *Abstract and Applied Analysis*, Volume **2011**, Article ID 649248, 10 pages.

[22] S. Araci, M. Acikgoz and J. J. Seo, Explicit formulas involving q-Euler numbers and polynomials, *Abstract and Applied Analysis*, Volume **2012**, Article ID 298531, 11 pages.

[23] S. Araci, D. Erdal, Higher order Genocchi, Euler polynomials associated with q-Bernstein type polynomials, *Honam Math. J.* 33(**2011**), no. 2, 173-179.

[24] S. Araci, X. Kong, M. Acikgoz, E. Şen, A new approach to multivariate q-Euler polynomials by using umbral calculus, arXiv:1211.4062 [math.NT].

[25] S. Araci, M. Acikgoz and A. Gürsul, Analytic continuation of weighted q-Genocchi numbers and polynomials, *Communications of the Korean Mathematical Society* (accepted for publication).

[26] S. Araci and M. Acikgoz, A note on the Frobenius-Euler numbers and polynomials associated with Bernstein polynomials, *Advanced Studies in Contemporary Mathematics* 22 (**2012**), no. 3, 399-406.

[27] S. Araci, M. Acikgoz, H. Jolany, J. J. Seo, A unified generating function of the q-Genocchi polynomials with their interpolation functions, *Proc. Jangjeon Math. Soc.* 15 (**2012**), no. 2, 227–233.

[28] S. Araci, M. Acikgoz, K. H. Park and H. Jolany, On the unification of two families of multiple twisted type polynomials by using p-adic q-integral on $\mathbb{Z}_p$ at $q = -1$, *Bulletin of the Malaysian Mathematical Sciences and Society* (In press).

[29] S. Araci, J. J. Seo and D. Erdal, New construction weighted (h, q)-Genocchi numbers and polynomials related to Zeta-type functions, *Discrete Dynamics in Nature and Society*, Volume 2011 (**2011**), Article ID 487490, 7 pages.

[30] Yuan He and Chunping Wang, Some formulae of products of the Apostol-Bernoulli and Apostol-Euler polynomials, *Discrete Dynamics in Nature and Society*, vol. 2012 (**2012**), Article ID 927953, 11 pages.

[31] S. Roman The umbral calculus, *Dover Publ. Inc. New York*, **2005**.